## Comparing the Adomian method and Adomian-Duan-Rach method for the solutions of a fourth-order nonlinear differential equation with four specified boundary values

## Navid Bozorgan<sup>1,\*</sup>, Mohsen Talebzadegan<sup>2</sup>

- <sup>1</sup> Department of Mechanical Engineering, Abadan Branch, Islamic Azad University, Abadan, Iran
- <sup>2</sup> Department of Mechanical Engineering, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran

Received: April 2016, Revised: May 2016, Accept: July 2016

## **Abstract**

In this paper, a fourth-order nonlinear differential equation with four specified boundary values is solved by the modified Adomian-Duan-Rach decomposition method. This modification also avoids solving a sequence of nonlinear algebraic equations for the undetermined coefficients fraught with multiple roots, thus more rapid rate of convergence observed for the Adomian decomposition series. In this method, all of the boundary values utilized before determining the Adomian polynomials. Example presented in this study showed that this method can be accelerated the convergence of the Adomian series with high accuracy for solving boundary value problems for higher order nonlinear differential equations.

\*Corresponding author: n.bozorgan@gmail.com

**Keywords:** Adomian decomposition method, Nonlinear differential equations, Boundary value problems.

Please cite this article as:

N. Bozorgan, M. Talebzadegan, Comparing the Adomian method and Adomian-Duan-Rach method for the solutions of a fourth-order nonlinear differential equation with four specified boundary values, Journal of Energy Conversion, (1-2) (2011) 31-34 [In Persian]